TYPE 1 DIABETES KIT Insulin Dependent NAME:	Must have access to food at any and all times Vomiting: Life threatening.	TARGET BLOOD SUGAR RANGE:
DOB:	Notify the parents	LOWS (BLOOD SUGAR BELOW):
ADDRESS:	immediately!	Signs (one or multiple): Irritability,
	CONTENT LIST	confusion, shakiness, impaired coordination, slurred speech.
		Treatment: Give fast acting sugar (small juice box, glucose tablets, Smarties, Skittles, glucose).
ATTACH YOUR CHILD'S PHOTO HERE		IN THE CASE OF EXTREMELY LOW BLOOD SUGAR (POSSIBLY BELOW 40) CAUSING SEIZURES, INABILITY TO INGEST FOODS, OR UNRESPONSIVENESS, ADMINISTER GLU- CAGON (0.5 MG) AND CALL 9-1-1. THIS IS AN IMMEDIATE THREAT TO LIFE THAT COULD CAUSE BRAIN DAMAGE OR
CONTACTS Emergencies, call 9-1-1		DEATH.
MOTHER:		HIGHS (BLOOD SUGARS ABOVE):
() - <u>FATHER:</u> () -		Signs (one or multiple): Lethargy, crankiness, dry throat, thirst, frequent need to use the restroom.
HOSPITAL: DOCTOR: () -		Treatment: Needs insulin. Must have access to bathroom. Check for ketones. If left untreated, organ damage will occur. Notify parent(s).